



**UNIVERSITI PUTRA MALAYSIA**

**DETECTION AND IDENTIFICATION OF JAAGSIEKTE RETROVIRUS  
ASSOCIATED WITH SHEEP PULMONARY ADENOMATOSIS**

**ZHU ZE**

**FPV 2002 5**

**DETECTION AND IDENTIFICATION OF JAAGSIEKTE RETROVIRUS  
ASSOCIATED WITH SHEEP PULMONARY ADENOMATOSIS**

**By**

**ZHU ZE**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the Requirement for the Degree of  
Doctor of Philosophy**

**January 2002**



Abstract of the thesis submitted to the Senate of Universiti Putra Malaysia  
in fulfillment of the requirement for the Degree of Doctor of Philosophy

**DETECTION AND IDENTIFICATION OF JAAGSIEKTE RETROVIRUS  
ASSOCIATED WITH SHEEP PULMONARY ADENOMATOSIS**

**By**

**ZHU ZE**

**January 2002**

**Chairman: Prof. Dr. Mohd Zamri Saad**  
**Faculty: Veterinary Medicine**

Sheep pulmonary adenomatosis (SPA) or jaagsiekte is a neoplastic disease of sheep, caused by jaagsiekte retrovirus (JSRV) infection in the lungs. In Malaysia, the disease was first observed in 1993, involving mainly the purebred imported sheep and their crossbreeds, and a long time after the first importation of sheep from Australia into Malaysia in 1987.

The JSRV was found to transform the alveolar type II cells in the lungs of sheep to become neoplastic cuboidal cells arranged in acinar or papillary patterns. The neoplastic cells were found to cover the entire alveolar wall with numerous macrophages filling the affected alveoli particularly the alveoli surrounding the neoplastic area.

The affected lung tissue and lung fluid were successfully processed to partially purify the materials before the JSRV was successfully detected under the

electron microscopy from several local SPA-affected sheep, which showed clinical signs typical of sheep pulmonary adenomatosis. Following the detection of JSRV under the electron microscopy, the virus was later confirmed as the JSRV by using the reverse-transcription polymerase chain reaction (RT-PCR) technique at the viral nucleic acid level.

The study, therefore, confirmed that JSRV is present in Malaysia and thus, should be considered in the diagnosis of sheep diseases and in controlling the spread of contagious diseases such as sheep pulmonary adenomatosis.

Following the confirmation of the jaagsiekte retrovirus in Malaysia, a RT-PCR diagnostic method was developed based on the U3 gene of the retrovirus. The method was able to identify the presence of retrovirus in the peripheral blood of sheep with and without clinical signs of sheep pulmonary adenomatosis. Thus, the technique can be used to identify subclinically infected animals for culling.

Abstrak tesis yang dikemukakan kepada senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**PENGESANAN DAN PENGENAL-PASTIAN VIRUS JAAGSIEKTE  
RETROVIRUS PENYAKIT ADENOMATOSIS PULMONARI BIRI-BIRI**

**Oleh**

**ZHU ZE**

**Januari 2002**

**Pengerusi: Prof. Dr. Mohd Zamri Saad**  
**Fakulti: Perubatan Veterinar**

Adenomatosis pulmonari bebiri (SPA) atau Jaagsiekte ialah sejenis penyakit neoplasia pada biri-biri akibat jangkitan peparu oleh jaagsiekte retrovirus (JSRV). Penyakit ini mula ditemui di Malaysia dalam tahun 1993 apabila melibatkan biri-biri baka tulen yang diimport dan kacukannya. Walau bagaimanapun, penyakit ini dikesan setelah sekian lama pengimportan pertama dibuat dari Australia.

Kajian menunjukkan virus JSRV menukar sel alveolus jenis II dalam paru-paru biri-biri menjadi sel epitelium kuboidal yang disusun bercorak asinus atau papilari. Sel-sel neoplasia ini di dapati menyelaputi keseluruhan bahagian dinding alveolus sedangkan makrofaj pula membanjiri rongga alveolus, terutama sekali di bahagian keliling kawasan berneoplasia.

Tisu peparu yang terlibat dan cecair telah diproses untuk penulenan separa. Hasil daripada penulenan separa ini, JSRV telah berjaya dikesan di dalam sampel

tersebut melalui pemeriksaan mikroskopi elektron. Pengesanan ini telah dilakukan ke atas beberapa ekor biri-biri Malaysia yang dikatakan telah dijangkiti penyakit tersebut serta menunjukkan petanda klinikal yang jelas. Setelah virus berjaya dikesan, virus ini kemudiannya telah dikenalpasti sebagai JSRV melalui ujian menggunakan teknik tindakbalas rangkaian polimerase pentranskripsian terbalik (RT-PCR) ke atas asid nukleus virus.

Maka, hasil kajian ini telah mengesahkan bahawa JSRV memang terdapat di Malaysia dan sepatutnya diambil-kira dalam membuat diagnosis penyakit biri-biri dan dalam menangani sebaran penyakit berjangkit seperti penyakit adenomatosis pulmonari biri-biri.

Berikutan pengenalan-pastian jaagsiekte retrovirus di Malaysia, kaedah diagnosis mengguna tindakbalas rangkaian polimerase pentranskripsian terbalik (RT-PCR) telah dihasilkan berdasarkan gen U3 retrovirus. Kaedah ini boleh mengenal-pasti kehadiran retrovirus dalam sampel darah biri-biri sama ada yang menunjukkan petanda klinikal penyakit adenomatosis pulmonari biri-biri atau tidak. Maka, teknik ini boleh diguna untuk mengenal-pasti biri-biri yang dijangkiti secara subklinikal agar ia boleh ditakai.

## ACKNOWLEDGEMENTS

I am exceedingly grateful to the chairman of my supervisory committee, Professor Dr. Mohd. Zamri Saad for his advice, guidance and patience throughout the planning and execution of this study. The stimulating discussions and constant encouragement were of great help for the mass of this work. Likewise his sacrifices and perseverance will ever be remembered.

Sincere thanks are also due to Assoc. Prof. Dr. Mohd Azmi Lila and Assoc. Prof Dr. Jasni Sabri who provided advice and direction, helpful in discussion and enlightening comments that have improved this study.

I would like to express my deep appreciation and gratitude to the followings individuals that have been of great help in my study. They include Dr. Md. Sabri Yusof, Dr. Zhang Shisheng, Dr. Lan Ganqiu, Dr. Mohd Effendy Abd. Wahid and members of the Faculty of Veterinary Medicine at the Universiti Putra Malaysia for sharing their technical skills and their ever available assistance.

Finally, I wish to express my deepest appreciation to my father and mother for their love and care as well as providing me with their blessing in carrying out this study.

I certify that an Examination Committee met on 24<sup>th</sup> January 2002 to conduct the final examination of Zhu Ze on his Doctor of Philosophy thesis entitled “Detection and Identification of Jaagsiekte Retrovirus Associated with Sheep Pulmonary Adenomatosis ” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**HASSAN HJ. MOHD DAUD, Ph.D.**

Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Chairman)

**MOHD ZAMRI SAAD. Ph.D.**

Professor / Deputy Dean  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**MOHD AZMI MOHD LILA, Ph.D.**


Associated Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**JASNI SABRI, Ph.D.**

Associated Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**MIKE SHARP, Ph.D.**

Professor  
Moredun Research Institute,  
UK.  
(External Examiner)



**AINI IDERIS, Ph.D.**

Professor  
Dean, School of Graduate Studies  
Universiti Putra Malaysia

Date: **1 MAR 2002**



This thesis submitted to Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy.



---

**AINI IDERIS, Ph.D.**  
Professor  
Dean, School of Graduate Studies  
Universiti Putra Malaysia

Date:

**14 MAR 2002**

## DECLARATION

I hereby declare that this thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not previously or concurrently submitted for any other degree at Unvesiti Putra Malaysia or other institutions.

  
ZHU ZE

Date: 1 MAR 2002

## TABLE OF CONTENTS

ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL SHEETS	viii
DECLARATION FORM	x
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF PLATES	xvi
LIST OF ABBREVIATIONS	xviii

## CHAPTER

1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2 1 The Sheep Pulmonary Adenomatosis Disease	4
2 2 Occurrence	5
2 3 Clinico-pathology	6
2 4 Experimental Transmission	8
2 5 Evidence that Jaagsiekte Retrovirus Causes Sheep Pulmonary Adenomatosis	9
2 6 Isolated and Expression of Jaagsiekte Retrovirus	11
2 7 Morphology and Morphogenesis of Jaagsiekte Retrovirus	15
2 8 Exogenous and Endogenous Jaagsiekte Retrovirus	20
2 9 The Long Terminal Repeat (LTR) of Jaagsiekte Retrovirus	26
2 10 Phylogenetic Analysis and Evolution of Jaagsiekte Retroviruses	31
2 11 Antibodies of Jaagsiekte Retroviruses	33
2 12 Oncogenesis of Sheep Pulmonary Adenomatosis	34
2 12 1 Transduction	34
2 12 2 Cis-activation	35
2 12 3 Trans-activation	35
2 12 4 Other mechanisms	36
2 13 Sheep Pulmonary Adenomatosis Shares Certain Characteristics With Human Bronchiolo-alveolar Carcinoma (BAC)	36
2 14 Diagnostic Techniques for Sheep Pulmonary Adenomatosis	37
2 14 1 History and Clinical Signs	37
2 14 2 Histology and Histochemical Tests	37
2 14 3 Electron Microscopy	38
2 14 4 Molecular Methods RT-PCR-Sequencing System	38
2 14 4 1 Polymerase Chain Reaction (PCR)	38
2 14 4 2 Reverse Transcription Polymerase Chain Reaction	39
2 14 4 3 DNA Sequencing	40
2 15 Research Focus on Sheep Pulmonary Adenomatosis	42



	<b>Page</b>
3 TISSUE CHANGES ASSOCIATED WITH RETROVIRUS IN NATURALLY OCCURRING SHEEP PULMONARY ADENOMATOSIS	44
3 1 Introduction	44
3 2 Materials and Methods	45
3 2 1 Animals	45
3 2 2 Sample Processing	46
3 2 3 Electron Microscopic Study	46
3 3 Results	47
3 3 1 Gross changes	47
3 3 2 Histopathological changes	49
3 3 3 Ultrastructural changes	52
3 4 Discussion	57
4 DETECTION AND IDENTIFICATION OF A RETROVIRUS ASSOCIATED WITH SHEEP PULMONARY ADENOMATOSIS	62
4 1 Introduction	62
4 2 Materials and Methods	64
4 2 1 Sources and Preparation of Samples	64
4 2 2 Sample Collection	65
4 2 3 Isolation and Partial Purification of Jaagsiekte Retrovirus For Electron Microscopy	66
4 2 4 Negative Electron Microscopy	67
4 2 5 Thin Section Electron Microscopy	67
4 2 6 Purification Jaagsiekte Retrovirus for Reverse Transcription Polymerase Chain Reaction	68
4 2 7 Jaagsiekte Retrovirus RNA Extraction	68
4 2 8 Synthesis of cDNA	70
4 2 9 Reverse-transcription Polymerase Chain Reaction	71
4 2 10 Restriction Endonuclease of Reverse Transcription Polymerase Chain Reaction Products	72
4 2 11 Detection of Polymerase Chain Reaction Products	73
4 2 12 Sequencing of Polymerase Chain Reaction Products	73
4 3 Results	75
4 3 1 Morphology of Jaagsiekte Retrovirus Particles	75
4 3 2 Polymerase Chain Reaction	77
4 3 3 Polymerase Chain Reaction Product Sequence	78
4 4 Discussion	78
5 UTILIZING PRIMER IN THE U <sub>3</sub> REGION OF JSRV LTR ESTABLISHED AN EXOGENOUS VIRUS SPECIFIC HEMI-NESTED RT-PCR	84
5 1 Introduction	84
5 2 Materials and Methods	86
5 2 1 Preparation of sample	86
5 2 2 Jagsiekte Retrovirus-RNA Extraction	86
5 2 3 Synthesis of cDNA	88

	<b>Page</b>
5 2 4 Hemi Nested Reverse Transcription Polymerase Chain Reaction	89
5 2 5 Detection of Polymerase Chain Reaction Product	90
5 2 6 Vectors and Competent Cells for Cloning	91
5 2 7 Jaagsiekte Retrovirus U3 Hemi-nested Reverse Transcription Polymerase Chain Reaction Amplification	92
5 2 8 Extraction of the Jaagsiekte Retrovirus 133 bp DNA Fragments from 1 5% Agarose Gels	94
5 2 9 Ligated into pT-Adv Vector	95
5 2 10 Transformation	96
5 2 11 Growth of Bacterial Clony	97
5 2 12 Isolation and Purification of Plasmid DNA	98
5 2 13 Restriction Enzyme Analysis	99
5 2 14 Polymerase Chain Reaction of the Plasmid	100
5 2 15 Sequencing	101
5 3 Results	102
5 3 1 Jaagsiekte Retrovirus U <sub>3</sub> Hemi-nested Reverse Transcription Polymerase Chain Reaction	102
5 3 2 Transformation	103
5 3 3 Restriction Enzyme Analysis	103
5 3 4 Polymerase Chain Reaction of the Plasmid	103
5 3 5 Sequence Analysis	105
5 4 Discussion	105
 6 PREVALENCE OF SHEEP PULMONARY ADENOMATOSIS AS DETERMINED BY THE NEWLY DEVELOPED U3 HEMI-NESTED ONE-TUBE RT-PCR	 110
6 1 Introduction	110
6 2 Materials and Methods	111
6 2 1 Reverse Transcription Polymerase Chain Reaction Based Sheep Pulmonary Adenomatosis Test Kit	111
6 2 2 Stability of the Hemi-nested One Tube Reverse Transcription Polymerase Chain Reaction Sheep Pulmonary Adenomatosis Diagnosis Kit	112
6 2 3 Prevalence of SPA in Sheep Farms	116
6 3 Results	117
6 3 1 Reverse Transcription polymerase chain reaction	117
6 3 2 Prevalence of Sheep Pulmonary Adenomatosis	118
6 3 3 Incidence of Sheep Pulmonary Adenomatosis Following Reverse Transcription Polymerase Chain Reaction	119
6 4 Discussion	121
 7 GENERAL DISCUSSION AND CONCLUSION	 122
 REFERENCES	 135
APPENDICES	153
VITA	180

## LIST OF TABLES

Tables	Page
2.1 Distribution of jaagsiekte retrovirus in the blood and tissue samples of sheep classified according to their sheep pulmonary adenomatosis status.....	15
4.1 The polymerase chain reaction mixture.....	72
5.1 RNase H reverse transcription reaction mixture.....	88
5.2 U <sub>3</sub> hemi-nested reverse transcription polymerase chain reaction Primers I, II, III.....	89
5.3 U <sub>3</sub> hemi-nested reverse transcription polymerase chain reaction first reaction mixture.....	89
5.4 U <sub>3</sub> hemi-nested reverse transcription polymerase chain reaction nested reaction mixture.....	90
5.5 Ligation reacted mixture.....	96
5.6 <i>EcoR</i> I restriction enzyme digested reaction mixture.....	99
5.7 Nested polymerase chain reaction mixture.....	100
6.1 Incidence of sheep pulmonary adenomatosis following reverse transcription polymerase chain reaction kit in farm 1.....	119
6.2 Incidence of sheep pulmonary adenomatosis following reverse transcription polymerase chain reaction kit in farm 2.....	120



## LIST OF FIGURES

Figures	Page
2.1 Genomic structures of endogenous and exogenous the type D retroviruses of premature stop code are indicated by a vertical bar, and underlined by an asterisk. ....	30
2.2 Phylogenetic analysis of the sheep type D retroviruses .....	32
4.1 The polymerase chain reaction strategy and localization of primers employed in this study. The <i>gag</i> reverse transcription polymerase chain reaction employing primers P1 and P2 were amplified a product of 229 bp. Nucleotide positions refer to the jaagsiekte retrovirus sequence published by York (York et al., 1992).....	71
4.2 Jaagsiekte retrovirus <i>gag</i> reverse transcription polymerase chain reaction products sequence.....	78
5.1 Restriction map and multiple cloning site of pT-Adv. Unique restriction sites were in bold. Restriction sites with asterisks were present only in the MCS and used to excise the inserted reverse transcription polymerase chain reaction product.....	93
5.2 The list showed the jaagsiekte retrovirus U <sub>3</sub> hemi-nested reverse transcription polymerase chain reaction product sequence.....	105

## LIST OF PLATES

Plates		Page
3.1	Photograph of lung from local bred siamese long tail sheep with pulmonary adenomatosis. Note the diffuse consolidation of the cranioventral region and patchy consolidation of the more dorsal region, the lesions were found diffusely distributed throughout the entire anterior lung.....	48
3.2	Photograph of lung from local bred siamese long tail sheep with pulmonary adenomatosis. ....	48
3.3	Photomicrograph, The histological lesion was the presence of tall cuboidal and columnar neoplastic epithelial cells that found lining along of the alveolar wall in the affected lobes. [H&E stain; x 100].....	50
3.4	Photomicrograph, Cuboidal tumor epithelial cells (arrow) with extensive vacuolation of cytoplasm and acinar lumens. [H&E stain; x 100].....	50
3.5	Photomicrograph, Neoplastic nodule with a clear papillary pattern. Myxoid tissue contains cells with elongated or round nuclei are present in the interstitium of the neoplastic tissue. [Toluidine blue, x 200].....	51
3.6	Photomicrograph. Some neoplastic foci. Foci of neoplastic epithelial proliferation of SPA. The acini are irregular and size, and have papillary in foldings. The interstitium contains mononuclear cells and discrete nodule of myxomatous tissue. [Toluidine blue, x 200].....	51
3.7	Photomicrograph, Histology section of an affected kidney showing severely shrunken glomeruls [H&E stain; x 400].....	52
3.8	Electron-micrograph showing ultrastructural features of type II neoplastic cells. The cells are markedly large and tall, found resting on a basement membrane. The cells have numerous small but mostly empty cytoplasmic vacuoles or inclusion (arrow). [x 4,500].....	54
3.9	Electromicrograph. Electron micrograph of columnar epithelium. Cells contain dense lamellar bodies (LB) predominantly at their apices and non-membrane bound lucent areas that contain glycogen (G) irregularly scattered in the cytoplasm. [ X 6,000].....	54
3.10	Transmission electron-micrograph showing the Jaagsiekte retrovirus particles observed in affected sheep pulmonary adenomatosis tissue. The particles have a mean diameter of 90 nm with small electron-lucent center surround by 2 shells. The cores are small and condense to form a rounded nucleoid (bar represents 48 nm). [x 80,000].....	55



<b>Plates</b>		<b>Page</b>
3.11	Transmission electron-micrograph showing Jaagsiekte retrovirus particles observed as a cluster of 3 intracytoplasmic virus particles (arrow) in a tumor cell. [x 40,000].....	55
3.12	Transmission electron-micrograph showing the Jaagsiekte retrovirus particles as two intracytoplasmic virus particles observed in a tumour cell [x 40,000].....	56
3.13	Transmission electron-micrograph showing the Jaagsiekte retrovirus particles in the final stages of budding from an apical microvillus. [x 120,000].....	56
4.1	Electron micrograph showing the Jaagsiekte retrovirus particles observed in negative stain electron microscopy. All the particles were observed in single form and rounded shape. [ x 65,000].....	76
4.2	Electron micrograph showing the higher magnification of the Jaagsiekte retrovirus negatively stained with PTA after fixation in osmium tetroxide. The particles were observed in single form and round-shaped of approximately 90 nm in diameter. [ x 65,000].....	76
4.3	Electrophoresis of the <i>gag</i> RT-PCR product showing the 229 bp band of jaagsiekte retrovirus. Lanes 1 & 6: Molecular weight markers (100 bp DNA ladder); lane 2: Lung fluid; lane 3: Lung tumor; lane 4: Mediastinal lymph nodes.....	77
5.1	U <sub>3</sub> JSRV Hemi-nested RT-PCR. Lanes 1 & 5: Molecular weight markers (100 bp DNA Ladder); lane 2: purified JSRV virus from lung fluid showing the 133 bp band; lane 3: Blood sample showing the 133 bp band; lane 4: Negative control.....	102
5.2	<i>EcoRI</i> Enzyme Analysis of Recombinant Plasmid pT-Adv-JSRV. Lane 1, 100 bp DNA ladder; Lanes 2 and 4, A recombinant plasmid from a white colony after digestion with <i>EcoRI</i> ; Lane 3, negative control (water); Lane 5: lambda <i>EcoR I</i> + <i>Hind III</i> markers.....	104
5.3	PCR of the insert in the recombinant plasmid. Lane 1, DNA molecular weight marker (100 bp DNA ladder); Lanes 2 and 3, plasmid PCR of primer pair PI and PIII.....	104
6.1	Hemi-nested RT-PCR SPA diagnosis Kit test result. Lane 1, Molecular weight markers (100 bp DNA Ladder); Lane 2 and 5 Blood sample; Lane 6, Positive Control.....	118

## LIST OF ABBREVIATIONS AND SYMBOLS

<b>A</b>	
$\alpha$	alpha
Amp	ampocillin
ATP	adenosine triphosphate
<b>B</b>	
$\beta$	beta
bp	base pair
BLV	bovine leukemia virus
<b>C</b>	
C	cytosine/core
$^{\circ}\text{C}$	degrees centigrade
cm	centimeter
cpm	counts per minute
<b>D</b>	
DNA	deoxy ribonucleic acid
DTT	1,4-Dithiothreitol
dsDNA	double stranded DNA
dsRNA	double stranded RNA
<b>E</b>	
EDTA	ethylenediaminetetra-acetic acid
ELSA	enzyme-linked immunosorbent assay
EM	electron microscopy
<b>F</b>	
Fig.	Figure
<b>G</b>	
g	gram
GA	glutaraldehyde
<b>H</b>	
H	hour
H & E	haematoxylin and eosin staining
<b>I</b>	
Ig G	immunoglobulin G
IPTG	isopropyl- $\beta$ -D-thiogalactopyranoside



<b>J</b>	
JSRV	jaagsiekte retrovirus
<b>K</b>	
kb	kilobase
kg	kilograms
<b>L</b>	
L	liter
LB	Luria broth
<b>M</b>	
$\mu$	micro
$\mu\text{g}$	microgram ( $10^{-6}$ g)
$\mu\text{l}$	micro liter ( $10^{-6}$ L)
$\mu\text{M}$	micro molar ( $10^{-6}$ M)
M	Molar
mg	magnesium
min	minute
ml	millitre
mm	millimeter ( $10^{-3}$ L)
mmol	millimolar
MMTV	mouse mammary tumor virus
MPMV	Mason-Pfizer monkey virus
mRNA	messenger RNA
MuSV	mirine sarcoma virus
MVV	Maedi-Visna virus
<b>N</b>	
Na	sodium
ng	nanogram ( $10^{-9}$ g)
nmol	nanmolar
<b>O</b>	
OD	optical density
OH	hydroxyl radical
ORF	open reading frame
<b>P</b>	
P	polymerase protein
PBS	phosphate buffered saline
PCR	polymerase chain reaction
pH	Puissance hydrogene
pmole	picomole

<b>R</b>	
RBC	red blood cell
RNA	ribonucleic acid
RT	reverse transcriptase,
rpm	revolutions per minute
<b>S</b>	
s	second
SDS	sodium dodecyl sulfate
SDW	sterile distilled water
SMRV	squirrel monkey retrovirus
SPA	sheep pulmonary adenomatosis
ssDNA	single stranded DNA
ssRNA	single stranded RNA
<b>T</b>	
TBE	Tris-boric acid-EDTA buffer
TBS	tris-buffered saline
TEM	transmission electron microscopy
<b>U</b>	
U	unite
UPM	Universiti Putra Malaysia
UV	ultraviolet
<b>V</b>	
V	volt
Vol	volume
v/v	volume / volume
<b>W</b>	
w/v	weight / volume
<b>X</b>	
Xg	centrifugal force
X-Gal	5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside

## CHAPTER 1

### INTRODUCTION

Sheep pulmonary adenomatosis (SPA), ovine pulmonary carcinoma (OPC) and jaagsiekte are alternative terms describing a contagious lung neoplasm of sheep. It is caused by a jaagsiekte retrovirus (JSRV) that is transmitted by the aerosol route. The clinical signs include rapid breathing, moist rales, coughing and production of a pulmonary fluid that drains from the nostrils. The dyspnea is most evident when affected animals are being herded leading to the African term ‘jaagsiekte’ or driving sickness for the disease.

Sheep pulmonary adenomatosis is classified as a bronchioloalveolar carcinoma (Stunzi et al., 1974). It is a neoplastic disease of sheep in which the jaagsiekte retrovirus transforms the alveolar type II cells to become neoplastic cuboidal epithelial cells arranged in acinar or papillary patterns (Rosadio et al., 1988). The disease was first described in South Africa and since then in many parts of the world including Europe (Bassett et al., 1989; Houwers et al., 1984), the United States of America (Culip et al., 1982). In Malaysia, the disease was first observed in 1993 involving mainly the purebred imported sheep and their crossbreds (Zamri-Saad et al., 1996).

The first reported case in many countries has been associated with importation of susceptible sheep into the country (Houwers et al., 1984; Bassett et al., 1989; Krishnan et al., 1994), and disease incidence usually increased

dramatically to involve many heads of animal and more sheep farms (Hunter et al., 1984; Zamri-Saad et al., 1996).

Based on earlier studies on the etiology, clinical and histopathology of sheep pulmonary adenomatosis, studies reported in this thesis were conducted with the following hypotheses:

Jaagsiekte retrovirus is present in Malaysia, and it would be confirm by using electron microscopy and the reverse-transcription polymerase chain reaction (RT-PCR) technique at the viral nucleic acid level.

The genomic RNA of the jaagsiekte retrovirus contains a unique nucleotide sequence, which is not available in the genome of other organisms or the host cells. If such area can be defined, specific DNA primers could be used to amplify the target region in the virus-specific manner by the hemi-nested reverse transcription polymerase chain reaction. Thus will enable the development of a diagnostic tool.

Since the disease was fairly new in this country (Zamri-Saad et al., 1996), the disease prevalence, transmission and development were unknown. Thus the objectives of this study were:

1. To study the tissue changes caused by jaagsiekte retrovirus following naturally occurring sheep pulmonary adenomatosis.
2. To describe sheep pulmonary adenomatosis and jaagsiekte retrovirus characteristics by electron microscopy.

3. To detect and confirm the presence of jaagsiekte retrovirus from several case of naturally occurring sheep pulmonary adenomatosis.
- 4 To develop the hemi-nested reverse-transcription polymerase chain reaction for detection of the jaagsiekte retrovirus. Cloning and sequencing of the suitable jaagsiekte retrovirus gene.
5. To determine and statistical analysis of the prevalence of the jaagsiekte retrovirus associated sheep pulmonary adenomatosis in Malaysia.

## CHAPTER 2

### LITERATURE REVIEW

#### 2.1. The Disease

Sheep pulmonary adenomatosis (SPA), jaagsiekte and ovine pulmonary carcinoma (OPC) are alternative terms describing a contagious lung neoplasm of sheep, apparently caused by a retrovirus known as the jaagsiekte retrovirus (JSRV). The virus is a slow virus with long incubation period, transmitted from the carrier ewes to the lambs by aerosol. There are occasional metastases to regional lymph nodes and other organs. The clinical signs of sheep pulmonary adenomatosis include rapid breathing, moist rallies, coughing, and over-production of a pulmonary fluid that drains from the nostrils. The dyspnea is most evident when affected animals are being herded leading to the Afrikaans terms for the disease of jaagsiekte, or 'driving sickness'.

Sheep pulmonary adenomatosis was first described in South Africa and since then in many parts of the world including Europe, the United States of America and Africa. In Malaysia, the disease was first observed in 1993 involving mainly the purebred imported sheep and their crossbreed (Krishnan et al., 1994). The first reported case in many countries has always been associated with importation of susceptible sheep into the country (Krishnan et al., 1994). Following the first incidence, the disease incidence usually increases dramatically to involve many more heads of animal and more sheep farms (Shah et al., 1997).



Sheep pulmonary adenomatosis is classified as bronchiole-alveolar carcinoma (Stunzi et al., 1974) with morphological similarities to human bronchiole-alveolar carcinoma (BAC), which constitutes between 3-5% of human pulmonary neoplasm (Sing et al., 1981; Perk et al., 1982). Both are relatively well-differentiated tumors of alveolar type 2 or non-ciliated bronchiolar (Clara) cells that exhibit malignant growth characteristics and have a peak occurrence in adults. Furthermore, the etiologies of both sheep pulmonary adenomatosis and bronchiole-alveolar carcinoma have been associated with a genetic factor (Wu et al., 1988). Studies on naturally occurring retrovirus-induced neoplasm, primarily lymphomas, have contributed immensely to the understanding of this disease.

## **2.2. Occurrence**

Sheep pulmonary adenomatosis was first recognized in South Africa in the 19th century as a cause of dyspnea in sheep being driven. Hence the origin of the Afrikaans name 'jaagsiekte', meaning driving sickness (Verwoerd et al., 1985). Subsequent reports documented the occurrence of sheep pulmonary adenomatosis as a sporadic or endemic disease of sheep and occasionally goats (Sharma et al., 1975) in all continents of the world except Australia (Verwoerd et al., 1985). Sheep pulmonary adenomatosis has been recognized as a common disease of sheep in Peru (Cuba Caparo et al., 1961; Snyder et al., 1983) and Chile (Shultz et al., 1965). Based on gross examination of lungs, the rate of diagnosis of sheep pulmonary adenomatosis among male was 1.5 times more than that of female, but the differences in management of the sexes could not be excluded as a cause of the